

FIG. 1




FIG. 2




FIG. 3





FIG. 4



0997322-113001
FOOT-2226660

FIG. 5




FIG. 6




FIG. 7




FIG. 7A





FIG. 8



05957822.113001
FOOT 228/5650

FIG. 9




FIG. 10




FIG. 13

10

FIG. 11

12

46

46

48

50

50

10

FIG. 12

44

48



46

50

50

10

44



BEST AVAILABLE COPY

FIG. 14





FIG. 15



FOOT " 223/6660

BEST AVAILABLE COPY

FIG. 16





FIG. 17



FOOT 228/6560

BEST AVAILABLE COPY

FIG. 18





FIG. 19



FOOT" 228/6660

BEST AVAILABLE COPY

FIG. 20





FIG. 21



100E T " 22B / 6660

BEST AVAILABLE COPY

FIG. 22




FIG. 23



0997822-113001

FIG. 24




FIG. 25




FIG. 26

Levels of Acridine derivative in PRBC treated with 200 μ M
Acridine Compound and 2 mM GSH




FIG. 27

Total level of Glutathione (oxidized and reduced) in
PRBC treated with 200 μ M Acridine Compound and 2
mM GSH





FIG. 28

% Hemolysis in PRBCs treated with
200 μ M Acridine Compound and 2 mM GSH



0997222-113001